

# POLYSUBSTANCE USE FACTS

## WHAT IS POLYSUBSTANCE USE?

The use of more than one drug, also known as polysubstance use, is common. This includes when two or more are taken together or within a short time period, either intentionally or unintentionally.

Intentional polysubstance use occurs when a person takes a drug to increase or decrease the effects of a different drug or wants to experience the effects of the combination. Unintentional polysubstance use occurs when a person takes drugs that have been mixed or cut with other substances, like fentanyl, without their knowledge.


Whether intentional or not, mixing drugs is never safe because the effects from combining drugs may be stronger and more unpredictable than one drug alone, and even deadly.

### What about prescription drugs?

The dangers of polysubstance use also apply to prescription drugs. Always let your doctor know what drugs you are taking to prevent any adverse reactions with newly prescribed medications. Never take pills that did not come from a pharmacy and weren't prescribed to you.

## GET THE FACTS

---



50%

In 2019,  
nearly half of  
drug overdose  
deaths involved  
multiple drugs.<sup>2</sup>

# THE DANGERS OF POLYSUBSTANCE USE



## Mixing Stimulants

**Examples of stimulants:** ecstasy (MDMA), cocaine, methamphetamines, amphetamines (speed)

Stimulants (also known as uppers) can increase your heart rate and blood pressure to dangerous levels and increase your risk of several serious health problems.

Combining stimulants may even directly or indirectly increase your risk of: brain injury, liver damage, heart attack, stroke.



## Mixing Depressants

**Examples of depressants:** opioids (prescription opioids, heroin, morphine, oxycodone, hydrocodone, fentanyl), benzodiazepines

Depressants (also known as downers) can slow down your breathing and increase your risk of several adverse health outcomes.

Combining depressants can also directly or indirectly increase your risk of: damage to the brain and other organs, overdose, death.



## Mixing Stimulants and Depressants

Mixing stimulants and depressants doesn't balance or cancel them out. In fact, the results of combining drugs are unpredictable, often modifying or even masking the effects of one or both drugs. This may trick you into thinking that the drugs are not affecting you, making it easier to overdose.



## Drinking alcohol while using other drugs

Drinking alcohol while using other drugs isn't safe. Alcohol is a depressant with similar effects to other downers. Mixing alcohol with other drugs can increase your risk of overdose and serious damage to the brain, heart, and other organs.

# WHAT TO DO IF YOU THINK SOMEONE IS OVERDOSING

It may be hard to tell whether a person is high or experiencing an overdose. If you aren't sure, it's best to treat the situation like an overdose—you could save a life.

1

Call 911 immediately.\*

2

Administer naloxone,\*\* if available.

3

Try to keep the person awake and breathing.

4

Lay the person on their side to prevent choking.

5

Stay with them until emergency workers arrive.

1. NCHS, National Vital Statistics System. Estimates for 2020 are based on provisional data. Estimates for 2015-2019 are based on final data (available from: <https://www.cdc.gov/nchs/hvssa/vsrr/drug-overdose-deaths.html>).

2. O'Donnell J, Gladde RM, Mattson CL, Hunter CT, Davis NL. Vital Signs: Characteristics of Drug Overdose Deaths Involving Opioids and Stimulants - 24 States and the District of Columbia, January-June 2019. MMWR Morb Mortal Wkly Rep 2020; 69:1189-1197. DOI: <http://dx.doi.org/10.15585/mmwr.mm6935a1>

